**COV FR 14230** 





Promising qualities characterise this variety: size potential, attractivity of the fruits

Adapted for all production areas

This variety will benefit to be grafted on a **vigorous** rootstock

# **APOLA**

Certification: in progress clone Ctifl n°14 401

#### Maura® period

## (0) Description du fruit

The fruit has a round, very attractive shape, with a small dip at its top. The fruit skin colour varies from garnet red to pink, and is characterised by a low pubescence.


The semi-freestone is small to medium sized. Firmness of the fruit is good.

The flesh is creamy white, and characterised by a light red lining under the skin, which is not found around the stone. The flavor is good, the acidity level is mild. The general eating quality is good, the aromatic intensity very satisfying. The texture is smooth and juicy.

#### **Production**

In good production conditions, the yield can reach 40T/Ha

The dominant fruit size is AA, with an average weight of 180-190g.





#### Fruit at harvest time

Av.	LOW		1EDIUM	HIGH	
BRIX (%)*			10 12		
Acidity (meq / 100 g)*	6	8			
Aroma intensity			6	7	

\*Pimprenelle measures



#### Fruit characteristics

Split cores susceptibility: weak

Rot susceptibility: weak


Sensitivity to handling: low to medium



Information
on this
technical bulletin
01/2022
is indicative only
and strongly
depends
upon production
conditions,
harvest time,
and soil/climatic
conditions
of the year.

This is
a protected variety,
which requires
a license
and is the subject
of
royalties payments.

Propagation and production can only be done with IPS agreement.





### **Blooming**

Bud break: semi early

Period : semi early, Patty® + 4 days

• Bloom set: medium to high

• Fruit setting: low to medium


Thinning: weak

### **铅 Tree**

Habit: semi-erect to erect

Vigor: medium to strong

Wood qualities : good



- Shape pruning: an open structure must be encouraged. For that, summer topping must be done during the 1st and the 2nd years.
   Subsequently, the trees will be trained by pruning rather than bending.
- Regular pruning: 2 year old wood and the strongest flowering shoots are preserved, even if they are in an acrotone position on the branches.
- Summer pruning: Essential to clean the inner part of the trees.
   Pruning can be done after the harvest to favor new buds on old branches.

Productivity : strong

Fruit set : rapid

