



Yellow nectarine

# HONEY BLAZE<sup>COV</sup>

COV EU 16 996

5 days before Big Top<sup>®</sup>



Very nice presentation for this highly productive variety

- Big Top<sup>®</sup> period

- Very well adapted in all production areas


## 🕒 Fruit description

- ◆ **Shape** : circular, regular, with a little dip at the top
- ◆ **Skin** : very attractive shiny red. The fruit skin is thick, almost not punctuated.
- ◆ **Core** : medium sized, semi clingstone
- ◆ **Firmness** : very good
- ◆ **Flesh** : smooth and juicy
- ◆ **Organoleptic qualities** : very good , with a sweet flavour

## 🏭 Production

In good production conditions, the yield can reach **30-35T/Ha.**

The dominant fruit size is **A/AA**, with an average weight of **160-170 g.**



## 📦 Fruit at harvest time

|                               | LOW | MEDIUM | HIGH |
|-------------------------------|-----|--------|------|
| <b>BRIX (%)*</b>              |     | 10 12  |      |
| <b>Acidity (meq / 100 g)*</b> |     | 8 10   |      |
| <b>Aroma intensity</b>        |     |        | 6 7  |

\* Pimprenelle measures

## 🗨️ Fruit characteristics

- ◆ **Split cores susceptibility** : weak
- ◆ **Rot susceptibility** : weak
- ◆ **Sensitivity to handling** : weak



Information on this technical bulletin 01/2022 is indicative only and strongly depends upon production conditions, harvest time, and soil/climatic conditions of the year.

This is a protected variety, which requires a license and is the subject of royalties payments.

Propagation and production can only be done with IPS agreement.

## 🌸 Blooming

- ◆ **Bud break** : mid season
- ◆ **Period** : mid season
- ◆ **Bloom set**: strong to very strong
- ◆ **Fruit setting** : very strong
- ◆ **Thinning** : strong



## 🌳 Tree

- ◆ **Habit** : semi erect
- ◆ **Vigor** : strong
- ◆ **Wood qualities** : very good
- ◆ **Shape pruning** : topping must be done only the 1st year. The opening of the scaffold branches will take place naturally. The very basiton character of this variety dictates the pinching of the sub scaffold branches to favor the growth of the main scaffold branches.
- ◆ **Regular pruning** : on vigorous and high quality flowering shoots. Do not leave fruits on the last 1/3 of the scaffold branches to prevent the collapse of the branches.
- ◆ **Summer pruning** : indispensable because this variety produces numerous vigorous shoots. Post harvest pruning will very probably lead to the regrowth of the flowering shoots before winter.
- ◆ **Productivity** : strong
- ◆ **Fruit set** : very rapid



*riale Violette? L: 98.  
A 01/22*