

Yellow peach

ROYAL SUMMER®

COV EU 27 232

Certification : clone INFEL® n°9266

3 to 5 days after Royal Mona®



Very nice variety
with **high fruit
size and yield
potential**

Very attractive
fruit with nice
aromatics
flavours


① Fruit description

- ◆ **Shape** : circular, slightly dysimetric, with a space at the stem end
- ◆ **Skin** : intense dark red. Skin is thin and has very few fuzz.
- ◆ **Core** : big sized, semi clingstone
- ◆ **Firmness** : good
- ◆ **Flesh** : crunchy, smoothy and juicy
- ◆ **Organoleptic qualities** : good, smooth and aromatic

👁 Production

In good production conditions, the
yield can reach **35-40T/Ha.**

The dominant fruit size is **A/AA**, with
an average weight of **160 -170 g**



Fruit at harvest time

	LOW	MEDIUM	HIGH
BRIX (%)*		10 12	
Acidity (meq / 100 g)*	6 8		
Aroma intensity		6 7	

* Pimprenelle measures

💡 Fruit characteristics

- ◆ **Split cores susceptibility** : weak
- ◆ **Rot susceptibility** : weak
- ◆ **Sensitivity to handling** : weak

Information
on this
technical bulletin
01/2022
is indicative only
and strongly
depends
upon production
conditions,
harvest time,
and soil/climatic
conditions
of the year.

This is
a protected variety,
which requires
a license
and is the subject
of
royalties pay-
ments.

Propagation
and
production
can only be done
with IPS agree-
ment.

🌸 Blooming

- ◆ **Bud break** : semi early
- ◆ **Period** : semi early
- ◆ **Bloom set** : medium
- ◆ **Fruit setting** : medium
- ◆ **Thinning** : low

🌳 Tree

- ◆ **Habit** : semi erect to erect
- ◆ **Vigor** : strong
- ◆ **Wood quality** : good
- ◆ **Shape pruning** : topping must be done during the 1st and 2nd years. Subsequently the scaffold branches must be opened by bending and/or pruning. Be careful to select and maintain sub scaffold branches and flowering shoots for the following year.
- ◆ **Regular pruning** : on stiff flowering shoots and vigorous 2 year old wood
- ◆ **Summer pruning** : Indispensable to favor good quality bloom set and a nice fruit coloration with a proper light exposure. Post harvest pruning is recommended.
- ◆ **Productivity** : strong
- ◆ **Fruit set** : moderately rapid

