



The reference of early varieties

Seasoned flowering time

Low susceptibility to split cores

SUGAR TIME cov

COV EU 30 095

7 days before Royal Dixie®

(iii) Fruit description

Shape: circular to oval, quite regular

Skin: red intense. Skin is moderatly thick and has little fuzz.

Core: medium sized, semi clingstone

Firmness: good


Flesh: smooth and juicy

Organoleptic qualities: good, with smooth and aromatic flavours

Production

In good production conditions, the yield can reach 20-25T/Ha.

The dominant fruit size is A, with an average weight of 140 -150 g.





🚊 Fruit at harvest time

	LOW	MEDIUM	HIGH
BRIX (%)*	9	11	
Acidity (meq / 100 g)*	6 9		
Aroma intensity		5 6	

^{*} Pimprenelle measures

(1) Fruit characteristics

Split cores susceptibility: weak

Rot susceptibility: weak


Sensitivity to handling: weak



Information
on this
technical bulletin
01/2022
is indicative only
and strongly
depends
upon production
conditions,
harvest time,
and soil/climatic
conditions
of the year.

This is
a protected variety,
which requires
a license
and is the subject
of
royalties payments.

Propagation and production can only be done with IPS agreement.





Blooming

• Bud break : semi late

• **Period**: mid season

Bloom set: strong

Fruit setting: strong

Thinning: strong

铅 Tree

Habit: erect

Vigor : medium

Wood qualities : good

Shape pruning: topping must be done during the first and second years. Subsequently the scaffold branches will be opened by bending and/or pruning.

Regular pruning: on vigorous flowering shoots. Renewal branches
 will come by bending the vigorous 2 year old wood.

Summer pruning: Indispensable to take off the vigorous shoots,
 but be careful to not be excessively severe.

Productivity: very strong

Fruit set : rapid



